

INTERPRETING THE ARITHMETIC IN THOMPSON'S GROUP F

VALERY BARDAKOV[†] AND VLADIMIR TOLSTYKH

ABSTRACT. We prove that the elementary theory of Thompson's group F is hereditarily undecidable.

M. Sapir asked whether the elementary theory of Thompson's group F is decidable [9, a part of Question 4.16]. More specifically, he asked later whether F first-order interprets the restricted wreath product $\mathbf{Z} \wr \mathbf{Z}$. It is known that the elementary theory of the latter group first-order interprets the arithmetic, and is therefore hereditarily undecidable [8]. Recall that a first-order theory T is said to be *hereditarily undecidable* if every subtheory of T is undecidable.

The aim of the present note is to show that F indeed first-order interprets $\mathbf{Z} \wr \mathbf{Z}$, and hence the elementary theory of F is hereditarily undecidable.

Let J be any closed interval of $\mathbf{R}$ with dyadic endpoints. The group $\text{PL}_2(J)$ is the group of all continuous order-preserving piecewise linear maps from J into itself with finitely many breakpoints all occurring at dyadic rationals and with the slopes of their linear 'components' equal to powers of two. One of the standard realizations of Thompson's group F is given by the group $\text{PL}_2([0, 1])$ (for the background information on Thompson's group F the reader may consult [1, 2].) It is well-known that $\text{PL}_2([0, 1])$ is generated by the following maps x_0 and x_1 :

$$\alpha x_0 = \begin{cases} 2\alpha, & 0 \leq \alpha \leq \frac{1}{4}, \\ \alpha + \frac{1}{4}, & \frac{1}{4} \leq \alpha \leq \frac{1}{2}, \\ \frac{1}{2}\alpha + \frac{1}{2}, & \frac{1}{2} \leq \alpha \leq 1, \end{cases} \quad \alpha x_1 = \begin{cases} \alpha, & 0 \leq \alpha \leq \frac{1}{2}, \\ 2\alpha - \frac{1}{2}, & \frac{1}{2} \leq \alpha \leq \frac{5}{8}, \\ \alpha + \frac{1}{8}, & \frac{5}{8} \leq \alpha \leq \frac{3}{4}, \\ \frac{1}{2}\alpha + \frac{3}{4}, & \frac{3}{4} \leq \alpha \leq 1. \end{cases}$$

Note that we write maps of $\text{PL}_2([0, 1])$ on the right.

By the definition,

$$x_n = x_0^{-n+1} x_1 x_0^{n-1}$$

where $n \geq 1$ is a natural number.

For any map $f \in \text{PL}_2([0, 1])$ we define

$$\text{supp}(f) = (\text{the support of } f) = \{\alpha \in [0, 1] : \alpha f \neq \alpha\}.$$

Clearly,

$$\text{supp}(f^{-1}gf) = \text{supp}(g)f$$

for all $f, g \in \text{PL}_2([0, 1])$. It follows that if maps $f, g \in \text{PL}_2([0, 1])$ have disjoint supports, then they are commuting.

2000 *Mathematics Subject Classification.* 20F65 (03C35, 03C60).

[†]The first author gratefully acknowledges support from TÜBİTAK, the Scientific and Research Council of Turkey.

The restricted wreath product $\mathbf{Z} \wr \mathbf{Z}$ has the following presentation:

$$\mathbf{Z} \wr \mathbf{Z} \cong \langle a, b \mid \{[b, a^{-n}ba^n] : n \in \mathbf{Z}\} \rangle.$$

The group F is ‘riddled’ with copies of $\mathbf{Z} \wr \mathbf{Z}$, as, for instance, the results from [6] show. We shall demonstrate that one such a copy is given by the group G generated by $a = x_0^2$ and $b = x_1x_2^{-1}$ (cf. [3] where it is proved that the subgroup generated by x_0 and $x_1x_2x_1^{-2}$ is also isomorphic to $\mathbf{Z} \wr \mathbf{Z}$.)

Lemma 1. (i) *The supports S_k of the maps $a^{-k}ba^k = x_0^{-2k}x_1x_2^{-1}x_0^{2k}$ where $k \in \mathbf{Z}$ are pairwise disjoint open intervals such that*

$$\bigcup_{k \in \mathbf{Z}} \overline{S_k} = [0, 1],$$

where $\overline{J}$ is the closure of an interval J of $\mathbf{R}$;

(ii) *The group G is isomorphic to $\mathbf{Z} \wr \mathbf{Z}$.*

Proof. (i) A straightforward calculation shows that

$$\text{supp}(x_1x_2^{-1}) = \left(\frac{1}{2}, \frac{7}{8}\right) = \left(\frac{1}{2}, \frac{1}{2}x_0^2\right)$$

(to calculate the products of elements of $\text{PL}_2([0, 1])$ we suggest the reader use the method of the tree diagrams, see, for instance, [1, Section 1.2].)




FIGURE 1. Graphs of the maps $a^{-k}ba^k$ (the graph of b is $OABCZ$, the graph of $a^{-1}ba$ is $OCDEZ$, the graph of aba^{-1} is $OC'B'AZ$, etc.)

Hence

$$S_k = \text{supp}(a^{-k}ba^k) = \text{supp}(x_0^{-2k}x_1x_2^{-1}x_0^{2k}) = \left(\frac{1}{2}x_0^{2k}, \frac{1}{2}x_0^{2k+2}\right)$$

for all integer numbers k . Note that

$$\lim_{n \rightarrow +\infty} \frac{1}{2}x_0^{2n} = 1$$

and that

$$\lim_{n \rightarrow -\infty} \frac{1}{2}x_0^{2n} = 0.$$

Then

$$(2) \quad [0, 1] = \bigcup_{k \in \mathbf{Z}} \overline{S}_k = \bigcup_{k \in \mathbf{Z}} \left[\frac{1}{2}x_0^{2k}, \frac{1}{2}x_0^{2k+2} \right],$$

as required.

(ii) By (i), the maps $a^{-k}ba^k$ where k runs over $\mathbf{Z}$ generate a free abelian group of infinite rank. Hence every element g of $G = \langle a, b \rangle$ is uniquely written in the form

$$(2) \quad g = a^m \prod_{k \in \mathbf{Z}} a^{-k}b^{m_k}a^k$$

where in the latter product only finitely many terms are non-trivial. The element $b = x_1x_0^{-1}x_1^{-1}x_0$ is in the commutator subgroup F' of F . Then g is congruent to a^m modulo F' . As $F/F' = \langle x_0F', x_1F' \rangle$ is a free abelian group of rank two [1, 2], $m \neq 0$ implies that $g \neq 1$. If $m = 0$, then $g = 1$ if and only if all exponents m_k are zeroes. Thus the generators a, b satisfy exactly the relations that are corollaries of those given in the presentation of the group $\mathbf{Z} \text{ wr } \mathbf{Z}$ above. $\square$

Proposition 2. *The group G is first-order definable in $\text{PL}_2([0, 1])$ with the parameters x_0, x_1 .*

Proof. We are going to prove that the subgroup $\langle a \rangle$ generated by a and the subgroup $\langle a^{-k}ba^k : k \in \mathbf{Z} \rangle$ are both definable in $\text{PL}_2([0, 1])$ by means of first-order logic with the parameters x_0 and x_1 . This in view of the formula (2) will imply that $G = \langle a, b \rangle$ is $\{x_0, x_1\}$ -definable, completing the proof.

Let $J = [\eta, \mu]$ be a closed interval with dyadic endpoints. $\text{PL}_2^>(J)$ (resp. $\text{PL}_2^\geq(J)$) is the family $\{f\}$ of all maps in $\text{PL}_2(J)$ such that $\alpha f > \alpha$ (resp. $\alpha f \geq \alpha$) for all $\alpha \in (\eta, \mu)$. The following result is an immediate corollary of Lemma 2.16 in [7].

Lemma 3. *Let $z \in \text{PL}_2^>(J)$. Then for every $q = 2^m$ where $m \in \mathbf{Z}$ there is at most one map in $\text{PL}_2(J)$ commuting with z whose initial slope is q .*

By the initial slope of an f in $\text{PL}_2([\eta, \mu])$ we mean the right-sided limit at η of the derivative of f , or, less formally, the slope of the first linear ‘component’ of f .

Lemma 3 implies that the centralizer of x_0 in $\text{PL}_2([0, 1])$ is the subgroup $\langle x_0 \rangle$ generated by x_0 : indeed, the initial slope of x_0 is 2 and the initial slope of a power x_0^m of x_0 , obviously commuting with x_0 , is 2^m . The powers of $a = x_0^2$ are the squares of elements of $\langle x_0 \rangle$. Thus the family of powers of a is first-order definable in $\text{PL}_2([0, 1])$ with the parameter x_0 . It follows that the set $\{a^{-k}ba^k : k \in \mathbf{Z}\}$ is definable in F with the parameters x_0 and x_1 .

We claim that

$$\langle a^{-k}ba^k : k \in \mathbf{Z} \rangle = C(\{a^{-k}ba^k : k \in \mathbf{Z}\})$$

where $C(X)$ is the centralizer of a subset X of $\text{PL}_2([0, 1])$. If so, the subgroup $\langle a^{-k}ba^k : k \in \mathbf{Z} \rangle$ is $\{x_0, x_1\}$ -definable in $\text{PL}_2([0, 1])$, being the centralizer of an $\{x_0, x_1\}$ -definable set.

Claim 4. *Let $k \in \mathbf{Z}$. Then the map $g_k = a^{-k}ba^k|_{\overline{S}_k}$ where $S_k = \text{supp}(a^{-k}ba^k)$ is in $\text{PL}_2^{\geq}(\overline{S}_k)$ and its initial slope is 2.*

Proof. The semi-group $\text{PL}_2^{\geq}([0, 1])$ is preserved under conjugation by elements of $\text{PL}_2([0, 1])$. Consequently, $g|_S \in \text{PL}_2^{\geq}(\overline{S})$, where $S = \text{supp}(g)$, for any conjugate g of b , since $b = x_1x_2^{-1} \in \text{PL}_2^{\geq}([0, 1])$.

It is easily seen that both maps $a = x_0^2$ and $a^{-1} = x_0^{-2}$ are linear on the intervals $(0, 1/8)$ and $(7/8, 1)$ and hence differentiable on these intervals. Suppose that $g \in \text{PL}_2^{\geq}([0, 1])$ is differentiable at $\alpha \in (0, 1/8)$. Then, by the Chain Rule, the derivative of aga^{-1} at the point αa^{-1} exists and is equal to

$$(3) \quad \left. \frac{da^{-1}}{dt} \right|_{t=\alpha g} \cdot \left. \frac{dg}{dt} \right|_{t=\alpha} \cdot \left. \frac{da}{dt} \right|_{t=\alpha a^{-1}},$$

that is, to the derivative of g at α , provided that $\alpha g < 1/8$ and $\alpha a^{-1} < 1/8$.

By direct calculations we see that the second statement of the Claim is true for maps b, aba^{-1}, a^2ba^{-2} . The support $S_{-2} = (1/32, 1/8)$ of the map a^2ba^{-2} enters the interval $(0, 1/8)$. We prove the statement for all $k \leq -2$ by induction: by (3), for any $k \leq -2$, the initial slope of the map $a^{-k}ba^k|_{S_k}$ is equal to the initial slope of the map $a^{-k+1}ba^{k-1}|_{S_{k-1}}$. The argument for positive k is similar: we check directly that the map $a^{-1}ba$ whose support is contained in $(7/8, 1)$ behaves as described, and prove the statement for all $k \geq 1$ by induction. $\square$

Now let $f \in C(\{a^{-k}ba^k : k \in \mathbf{Z}\})$. Then for every $k \in \mathbf{Z}$ the restriction f_k of f on S_k commutes with g_k . By Claim 4 and Lemma 3, $f_k = g_k^{l_k}$ for a suitable integer l_k . Finally, part (i) of Lemma 1 implies that at most finitely many l_k are non-zero, whence

$$f = \prod_{k \in \mathbf{Z}} a^{-k}b^{l_k}a^k,$$

and the result follows. $\square$

Proposition 5. *The elementary theory of Thompson's group F is hereditarily undecidable.*

Proof. It is well-known that if a structure $\mathcal{M}$ first-order interprets with parameters a structure $\mathcal{N}$ whose elementary theory is hereditarily undecidable, then the elementary theory of $\mathcal{M}$ is too hereditarily undecidable [5].

According to [8], if a finitely generated almost solvable group S is not almost abelian, then S first-order interprets the ring of integers $\mathbf{Z}$ (see also [4].) It is easy to see that the group $\mathbf{Z} \text{ wr } \mathbf{Z}$ is solvable; in order to see that it is not almost abelian (that is, does not have normal abelian subgroups of finite index) one can work with the isomorphic group $G = \langle a, b \rangle$ introduced above. Suppose that φ is a surjective homomorphism from G onto a finite group of order n . Then

$$1 = \varphi(a)^n = \varphi(b)^n = \varphi(a^n) = \varphi(b^n).$$

The elements a^n, b^n are not commuting, and hence the kernel of φ is not abelian.

Thus the elementary theory of the group $\mathbf{Z} \text{ wr } \mathbf{Z}$ is hereditarily undecidable, for the elementary theory of the ring of integers is hereditarily undecidable [5, 10].

Now, by Proposition 2, Thompson's group F first-order interprets with parameters the group $\mathbf{Z}$ wr $\mathbf{Z}$, and we are done. $\square$

The authors would like to express their gratitude to Oleg Belegradek for interesting discussions and to thank Victor Guba and Mark Sapir for helpful information.

REFERENCES

- [1] J. Belk, Thompson's group F , Ph. D. Thesis, Cornell Univ., 2004, available at <http://www.math.cornell.edu/~belk/Thesis.pdf>.
- [2] J. Cannon, W. Floyd, and W. Parry, Introductory notes to Richard Thompson's groups, *Enseign. Math.* 42 (1996) 215–256.
- [3] S. Cleary, Distortion of wreath products in some finitely presented groups, 2005, preprint, [arXiv:math.GR/0505288](https://arxiv.org/abs/math/0505288).
- [4] F. Delon, P. Simonetta, Undecidable wreath products and skew power series fields, *J. Symb. Logic* 63 (1998) 237–246.
- [5] Yu. Ershov, Decision problems and constructivizable models, Nauka, Moscow, 1980.
- [6] V. Guba, M. Sapir, On subgroups of the R. Thompson group F and other diagram groups, *Mat. Sb.* 190 (1999) 3–60.
- [7] M. Kassabov, F. Matucci, Simultaneous conjugacy problem in Thompson's group F , 2006, preprint, [arXiv:math.GR/0607167](https://arxiv.org/abs/math/0607167).
- [8] G. Noskov, The elementary theory of a finitely generated almost solvable group, *Izv. Akad. Nauk SSSR Ser. Mat.* 47 (1983) 498–517.
- [9] Problem list of the workshop 'Thompson's Group at 40 Years' (Jan. 11-14 2004, American Institute of Mathematics, Palo Alto, California), available at <http://aimath.org/WWN/thompsonsgroup/thompsonsgroup.pdf>
- [10] R. Robinson, Undecidable rings, *Trans. Amer. Math. Soc.* 70 (1951) 137–159.

VALERY BARDAKOV, INSTITUTE OF MATHEMATICS, SIBERIAN BRANCH RUSSIAN ACADEMY OF SCIENCE, 630090 NOVOSIBIRSK, RUSSIA

E-mail address: `bardakov@math.nsc.ru`

VLADIMIR TOLSTYKH, DEPARTMENT OF MATHEMATICS, YEDITEPE UNIVERSITY, 34755 KAYIŞDAĞI, ISTANBUL, TURKEY

E-mail address: `vtolstyk@yeditepe.edu.tr`